

4.20; tail, 2.70; culmen, .45; tarsus, .75; middle toe, .70. *Hab.* New Granada to Guiana.

b. Markings on the crown in form of longitudinal streaks, or lines.

5. *G. FERRUGINEUM.* Tail dark brown, crossed by 7-9 continuous bands of bright rufous, about the same width as the brown ones. Above, varying from grayish brown to bright ferruginous, without lighter transverse bars on the dorsal region; beneath, varying from pure white to pale rufous, the stripes on the sides grayish brown, or rufous. Sides of the breast never speckled with lighter. Wing, 3.70-4.15; tail, 2.20-2.90; culmen, .45-.50; tarsus, .70-.80; middle toe, .70-.75. *Hab.* Tropical America, from southern border of U. S., (Arizona, Bendire; and probably Texas) to southern Brazil. Both coasts of Middle America, but apparently only the Atlantic slope of South America.

6. *G. INFUSCATUM.* Tail dark brown, crossed by 6-7 non-continuous bands of white, narrower than the dark ones. Above, varying from grayish brown to reddish umber and sepia (apparently never rufous). Beneath white, the stripes on the sides grayish brown, or dark brown. Sides of the breast not speckled.

Above, dark sepia, or blackish brown. Tail brownish black, or deep black. Wing, 3.70-3.90; tail, 2.50-2.90; culmen, .45; tarsus .65-.80; middle toe, .65-.70. *Hab.* Eastern South America (Paraguay to New Granada).

. . . . . var. *infuscatum.*

Above, grayish, or reddish, umber. Tail clear dark brown, or grayish umber. Wing, 3.60-3.90; tail, 2.35-2.75; culmen, .45-.50; tarsus, .65-.80; middle toe, .60-.70. *Hab.* Whole of Middle America, from the Rio Grande (perhaps in Texas) to Panama. . . var. *gnoma.*

7. *G. NANUM.* Tail brown, crossed by 8-12 continuous bands of reddish white, or rufous, not more than half as wide as the brown ones. Brown of the sides much broken anteriorly by white spotting, and forming