

nigris : loris et supercilio lato usque ad latera colli producto cinereis. Long. tot. 6·1 poll., culm. 0·7, alæ 3·0, caudæ 2·2, tarsi 1·1.

*Hab.* Efulen, Cameroons (*G. W. Bates*). Como River, 60 miles from Gaboon (*G. W. Bates*). R. Congo (*Petit : Mus. Brit.*).

*Obs.* A species easily distinguished from all the other African members of the genus by its black head and grey eyebrow.

## 2. TURDINUS CERVINIVENTRIS, n. sp.

*Turdinus fulvescens*, partim, auct. recent. (nec Cass.).

♂. Similis *T. fulvescenti*, Cass., sed corpore subtus cervino-brunneo, abdomine vix pallidiore : gula albida, griseo obscure striata : subcaudalibus cervinis, vix rufescentibus. Long. tot. 6·7 poll., culm. 0·8, alæ 2·95, caudæ 2·6, tarsi 0·95.

*Hab.* Conde, Congo (*Petit*), Chissambo (*Petit*), Yambuya, Aruwimi River (*Jameson*), Como River, Gaboon (*G. L. Bates*), Jaunde, Camaroons (*Zenker*).

*Obs.* The true *T. fulvescens*, Cass., of which the Museum contains two co-types, is a white-bellied species. Cassin appears to have confused two distinct species under the heading of *T. fulvescens*, as he observes : " In some specimens the entire underparts are dull reddish fulvous." These would be the new *T. cerviniventris*.

## 3. SCOPS HOLERYTHRA, n. sp.

Similis *S. icterorhynchæ*, Shelley, sed ubique cinnamomea, maculis albis scapularibus nullis, tectricibus alarum quoque albo haud maculatis : subtus læte cinnamomea, maculis longitudinalibus albis notata, sed minime fusco-vermiculata. Long. tot. 8·5 poll., culm. 0·8, alæ 5·5, caudæ 2·9, tarsi 1·05.

*Hab.* Efulen, Kamerun.

## 4. HAPALODERMA ÆQUATORIALE, n. sp.

Similis *H. narinæ*, sed minor : alæ fasciis vermiculatis nigris fortissime indicatis. Long. tot. 10·5 poll., culm. 0·8, alæ 4·7, caudæ 5·8, tarsi 0·65.

*Hab.* Efulen, Kamerun.