

possibly during the latter half of the Pleistocene, when land masses extended far beyond their present boundaries (Hope & Hope 1976: 30, Kikkawa *et al.* 1981: 1698). The possibility of such a chance transoceanic crossing by *Tyto novaehollandiae* from northern Australia to colonize islands in the Banda Sea by way of the Tanimbar Islands could be considered unrealistic were it not for additional evidence, that other avian species have in the past (e.g. *Cacatua pastinator goffini*, *Geopelia maugeus*) and at the present day do still follow this route (Hartert 1901, Schodde & Mathews 1977), and, accordingly, that considerable avifaunal interchange has taken place in the past between the southern Moluccas (Buru) and the Tanimbar Islands (see Hartert (1901) for similarities between a number of species and subspecies from these islands).

However, *T.n. cayelii* of Buru does show some similarity to *T. inexpectata* of Sulawesi in having the white dorsal spots replaced by brownish ones. Geographical position (see Audley-Charles (1981) for discussion on past land connections between Celebes and Australo-Papua) and morphological similarity of *T. inexpectata* to *T.n. cayelii* suggests that *inexpectata* may have budded off much earlier and speciated from the *T. novaehollandiae* line.

The origin of *manusi* is more difficult to explain but it may have followed a northeasterly route from Buru around northern New Guinea via the Moluccas, to the Admiralty Islands (see Salomonsen (1976: 595-6) for a possible route). Rothschild & Hartert (1914) also suggest that *manusi* is nearest to *cayelii*. *T.n. manusi*, in my opinion, on the other hand, exhibits little or no morphological similarity to *T. aurantia*, from New Britain, although *T. aurantia* presumably originated and diverged from Masked Owl stock (Schodde & Mason 1981: 77).

If *T. novaehollandiae* did follow this suggested evolutionary route, it is possible that there are undiscovered populations on some of the islands around the Moluccas and the Vogelkop. There are a number of publications discussing collections from intervening locations throughout this region without reference to *T. novaehollandiae*, while few specimens of the races discussed above exist in collections. K ln, for example (Hartert 1901), whilst collecting on Timorlaut (Tanimbar Islands), did not procure any specimens of *sorocula*, and the paucity of specimens from islands north of Australia may suggest a number of possibilities, e.g. a lack of systematic collecting of night birds, the inaccessibility of likely locations, that populations of these subspecies may be low in number or that they have a very patchy distribution — or a combination of all these factors.

***Tyto novaehollandiae calabyi* subsp. nov.**

Holotype. (Rijksmuseum van Natuurlijke Historie, Leiden) Reg. No. 42474, Field No. 305, adult ♂, Merauke (southern New Guinea—see Fig. 1), 22 April 1960, coll. A. Hoogerwerf. Measurements (mm) wing 305, tail 130, culmen (from cere) 24.3, tarsus 70.

Paratypes. (Rijksmuseum van Natuurlijke Historie, Leiden) Reg. No. 42475, ♀; (American Museum of Natural History) Reg. Nos. 425939, 425940—♂ and ♀ respectively.

Series studied. 2 ♂♂, 2 ♀♀; ♂, ♀ immatures.

Range. *T.n. calabyi* inhabits southern New Guinea, in the eucalypt savannahs between Merauke and the Fly River Delta (Fig. 1).

Diagnosis. *T.n. calabyi* differs from tropical island populations (*melwillensis*,

cayelii, *sorocula* and *manusi*) in its large size and its coarser spotting and speckling of the dorsal surface. It is distinguished from *galei* by its darker dusky and tawny, and less coarsely mottled, dorsal plumage (Plate 2), and larger size (Table 1).

Although *calabyi* of southern New Guinea is said not to differ from *novaeollandiae* (Mees 1964, 1982), *calabyi* can be distinguished by its consistently darker dorsal plumage (4 adults, 2 immatures) when compared with 7 medium and 4 dark phase birds from the northern limits of *novaeollandiae*'s range. To date there are no dark phase birds with an extensive russet wash to the dorsal surface recorded from southern New Guinea. Moreover, in differing from *novaeollandiae*, *calabyi* exhibits a number of characters similar to *galei*: pale feather pattern to the lesser wing coverts, secondary and primary flight feathers (contrasting with the rest of the dark dorsal plumage), lack of feathering and hair to the lower portion of the tarsus and toes, and proportionally larger tarsi and smaller tails (Table 1). Although New Guinean birds are as large as nominate *novaeollandiae*, they differ in their larger culmen and tarsus and their appendage ratios (Table 1).

Etymology. This subspecies is named after Dr. John H. Calaby in acknowledgment of his support and encouragement to my work over the years and because of his deep past interest in this species.

Acknowledgments. I would like to thank the curators of the Australian museums and the American Museum of Natural History for placing their collections of this species at my disposal, to express my gratitude to Dr. G. F. Mees (Rijksmuseum van Natuurlijke Historie, Leiden) for comments and material loaned, to Dr. R. Schodde for comments on the manuscript and arranging loans, to F. H. J. Crome for comments on the manuscript, and to Messrs. F. D. Knight for drawing the figure and J. Wombey who photographed the specimens.

References:

- Audley-Charles, M. G. 1981. 4. Geological History of the Region of Wallace's Line. In *Wallace's Line and Plate Tectonics* (T. C. Whitmore, Ed.): 24-35. Clarendon Press: Oxford.
- Hartert, E. 1901. On a collection of birds from the Timorlaut Islands. *Novit. Zool.* 8: 163-176.
- Hope, J. H. & Hope, G. S. 1976. Palaeoenvironments for man in New Guinea. In *The Origin of the Australians* (R. L. Kirk & A. G. Thorne, Eds.): 29-54. Australian Institute of Aboriginal Studies: Canberra.
- Kikkawa, J., Monteith, G. B. & Ingram, G. 1981. Cape York Peninsula: a major region of faunal interchange. In *Ecological Biogeography of Australia* (A. Keast, Ed.). *Mono. Biol.* 41: 1697-1742. The Hague: Dr. W. Junk.
- Mayr, E. 1941. *List of New Guinea Birds*. New York: Amer. Mus. Nat. Hist.
- Mees, G. F. 1964. A revision of the Australian Owls (Strigidae and Tytonidae). *Zool. Verhandl.* 65: 1-62.
- 1982. Birds from the lowlands of southern New Guinea (Merauke and Koembe). *Zool. Verhandl.* 191: 1-188.
- Rand, A. L. 1942. Results of the Archbold Expeditions. Birds of the 1936-37 New Guinea Expedition. *Bull. Amer. Mus. Nat. Hist.* 79: 289-366.
- Rand, A. L. & Gilliard, E. T. 1967. *Handbook of New Guinea Birds*. London: Weidenfeld and Nicolson.
- Rothschild, W. & Hartert, E. 1914. The birds of the Admiralty Islands, north of German New Guinea. *Novit. Zool.* 21: 282-292.
- Salomonsen, F. 1976. The main problems concerning avian evolution on islands. In *Proc. XVI Int. Orn. Congr.*: 585-602. Australian Academy of Science: Canberra.
- Schodde, R. & Calaby, J. H. 1972. The biogeography of the Australo-Papuan bird and mammal faunas in relation to Torres Strait. In *Bridge and Barrier: the natural and cultural history of Torres Strait* (D. Walker, Ed.): 257-300. Publ. BG/3. Dept. of Biogeogr. and Geomorphol.: Aust. Nat. Univ.

Schodde, R. & Mathews, S. J. 1977. Contributions to Papuan ornithology, V. *Survey of the Birds of Taam Island, Kai Group*. CSIRO Division of Wildlife Research Tech. Paper No. 33: 1-29.

Schodde, R. & Mason, I. J. 1980. *Nocturnal Birds of Australia*. Melbourne: Lansdowne Editions.

Address. Ian J. Mason, CSIRO Division of Wildlife and Rangelands Research, P.O. Box 780, Atherton, Queensland 4883, Australia.

© British Ornithologists' Club 1983

The nomenclature of *Buteo oreophilus*

by A. H. James and J. Wattel

Received 3 March 1983

In 1914, Hartert & Neumann described a species of buzzard from Koritscha, Ethiopia, under the name *Buteo oreophilus*, giving as the range of this species the mountain areas of central, northeast, and east Africa. Sclater (1919) was the first to suggest that the resident buzzards of southern Africa might also belong to the same species (see also Roberts 1940, Courtenay-Latimer 1941). It was not until 1957 that Rudebeck formally described and named the South African segment of this resident buzzard as *Buteo buteo trizonatus*, type locality Knysna, Cape Province. Hesitantly, he followed Meinertzhagen (1951) in considering both *oreophilus* and *trizonatus* as conspecific with the Eurasian *Buteo buteo*. Since then the taxonomic affinity of these forms has been examined by many authors, resulting in widely varying conclusions. A. H. J. intends to discuss the taxonomy of the group (James in prep.), but before doing so there is a need to clear up the nomenclatural confusion that has arisen recently from Brooke's (1974) conclusion that in his opinion Rudebeck (1957) was not the first author to describe the South African form of *Buteo*.

In 1830, Andrew Smith started publication of a series of papers in the South African Quarterly Journal, which he apparently intended should form a kind of catalogue of South African birds. In trying to identify these birds he referred to Levaillant's *Histoire Naturelle des Oiseaux d'Afrique* (1799-1808) and to Daudin's *Traité élémentaire et complet d'ornithologie* (1800). The latter work gave binominal scientific names. After citing these, Smith gave careful descriptions of the material he had at hand. At the time these were not meant to constitute descriptions of new species (Mees 1967, Clancey *et al.* 1971), but they were later considered as such by him and also by other authors.

Among the species of *Buteo* found in South Africa, Smith (1830) listed *Buteo tachardus* (*Falco tachardus* Daudin, 1800) and *Buteo desertorum* (*Falco desertorum* Daudin, 1800). *Falco tachardus* Daudin is based upon Le Tachard as depicted on Plate 19 in Levaillant (published in 1799), and *Falco desertorum* Daudin on Le Rougri (Plate 17 in Levaillant 1799). Both names, and particularly *F. desertorum* have later been applied to *Buteo buteo vulpinus* Gloger, 1833, a migrant to South Africa. However, Hartert (1914:1125) was of the opinion that Le Rougri is unidentifiable [. . . Figment of fantasy! The fox-red upperparts, the description of the underparts, the beautiful yellow bill, especially mentioned in the description, void the use of the name *desertorum*. On top of all this, the miraculous animal should be resident in South Africa and breed there]; and that Le Tachard refers to *Pernis apivorus* (Linnaeus,