

Male much darker on the underside than male of *T. s. maculosa* and of *T. s. beccarii*.

Wing, female 79–85, male 74–78 mm.

Distribution.—Peleng Island.

Type.—Female, Peleng Island, 20. vii. 1938 ; J. J. Menden leg.

Remarks.—Four ad. females, one semi-ad. female, six ad. males, were compared with several *T. s. maculosa*, *T. s. beccarii*, and *T. s. saturata* in the Berlin Museum, and with all specimens of available races by Mr. Kinnear in London, and by Dr. Junge in Leiden. Named in honour of Mr. N. B. Kinnear.

A more detailed description of the specimens will be published later on.

***Tyto rosenbergi pelengensis*, subsp. nov.**

Description.—Ad. male. Very similar to *Tyto rosenbergi rosenbergi* (Schlegel) from Celebes, but the feathers of the lower throat and upper breast with subterminal broken up black edges, thus giving these feathers a somewhat scaled appearance. Much smaller than *T. r. rosenbergi*.

Wing 296 mm., as compared with 335–360 mm. in *T. r. rosenbergi* (*vide* Meyer and Wilesworth, 'Birds of Celebes,' i. p. 110).

Distribution.—Peleng (Peling) Island.

Type.—Male, Peleng, 22. viii. 1938 ; J. J. Menden leg.

Remarks.—Only this one specimen was collected.

***Caprimulgus macrurus jungei*, subsp. nov.**

Description.—Male and female very similar to *Caprimulgus macrurus celebensis* Ogilvie Grant from Celebes, but the extension of the white tips of the two outer pairs of tail-feathers still more reduced, 15–17 mm., as compared to 25 mm. in *C. m. celebensis* and about 50 mm. in *C. m. macrurus*.

Wing, male 168–183 mm. Iris brown.

Distribution.—Taliaboe.

Type.—Male, Taliaboe, 13. x. 38 ; J. J. Menden leg.

Remarks.—Menden collected ten specimens, of which six are marked male and four female. If these indications are